
Health




HEALTH NEEDS OF COURT-INVOLVED YOUTH



Over **two-thirds** of youth in the justice system report health needs, including injuries, illness, and other medical problems.

JUVENILE OFFENDER INVOLVEMENT WITH CHILD WELFARE



60% of first-time juvenile offenders had a history of child welfare involvement.





Youth in the juvenile justice system are a **medically underserved** population. Health risks of youth in the juvenile justice system include:

- chronic health problems
- substance use
- risky sexual behavior

Many youth who are justice-involved report co-occurring health risks, between mental health disorders, sexually transmitted infections (STIs) and other sexrelated health complications, and higher rates of lifetime substance abuse.¹ Often, these health needs are not addressed until youth become involved with the justice system.

Child Maltreatment

Child maltreatment includes physical abuse, sexual abuse, psychological abuse, and neglect. Youth involved with courts suffer from high rates of neglect and abuse, both of which have been found to be strong predictors of violence and system involvement in adolescents.²

Neglect is the most common form of maltreatment; research shows that the quality of monitoring of a child by a parent or guardian is the strongest predictor of juvenile delinquency.³ Chronic neglect has been linked to other risk factors such as poverty, persistent substance abuse, mental health issues, and developmental disabilities.⁴

Experiencing maltreatment early in life puts youth at risk for post-traumatic stress symptoms, as well as for internalizing problematic behaviors, which in turn increases the likelihood of court involvement.⁵ One study found that youth who experienced maltreatment in childhood are 11% more likely to engage in delinquent behavior.⁶

Dual-Status Youth

Youth involved in both the child welfare and juvenile justice systems are referred to as **dual-status youth.** Youth with a history of child welfare involvement were referred for offenses **3 times as often** as youth with no child welfare involvement. Because childhood adversity disproportionately affects youth involved in the juvenile justice system, trauma-informed intervention programs should be developed to prevent juvenile delinquency and recedivism. 8

Chronic Health Problems

Court-involved youth present with high rates of traumatic injury, tuberculosis, dental problems, and sexually transmitted infections. Children who engage in delinquent and disruptive behaviors are at higher risk for poor physical health, especially resulting from injuries and accidents.⁹

Experiencing neglect alongside other adverse experiences in childhood is linked to health risks such as diabetes, heart disease, chronic obstructive pulmonary disease (COPD), cancer, and liver disease. See Adverse Childhood Experiences (ACEs) for more detail.

Health



Substance Use & Substance Use Disorders

Youth who are court involved are at high risk of developing substance use disorders, which signify a level of dependence that causes clinically significant impairment. Substance use disorders are in turn strongly associated with involvement in the juvenile justice system. Substance use puts youth at increased risk for mental health issues and is directly proportional to aggression and violent behavior. Research also shows that drug exposure interrupts brain development, which may contribute to impulsivity and risk-taking in adolescents. Acute substance use puts youth at higher risk of being identified by law enforcement and detained for drug and alcohol offenses, while chronic substance use increases the risk of violent offenses and recidivism. Substance use also increases the risk that adolescents will gravitate to peer contexts in which accidents and violence are more likely.

72% of youth who are court-involved develop a substance-use disorder.



In comparison, only 8% of youth in the general population develop a substance-use disorder.



Risky Sexual Behavior

Peer rejection, affiliation with deviant peers, and juvenile delinquency all increase the likelihood of risky sexual behavior both in adolescence and early adulthood. Peer rejection during childhood is especially a risk factor for girls engaging in risky sexual behavior. Risky sexual behavior, including having multiple sexual partners, initiating sexual activity at a young age, and not using contraception, increases the likelihood of health risks such as sexually transmitted infections and unintended pregnancy. A Risky sexual behavior in adolescents is also associated with sexual abuse, particularly among youth in the child welfare system.

One study found that girls with post-traumatic stress symptoms are more than **7 times more likely** to engage in unprotected sexual intercourse than girls without these symptoms.¹⁵





Another study found that depressive symptoms and drug use, both of which are found at heightened levels in court-involved youth, are related to a greater number of sexual partners in boys, and inconsistent use of protection in girls.¹⁶

All of these health risks are interrelated.

This highlights the importance of developing standards of care, as well as coordination between the justice system, health providers, and community providers to address the mental and behavioral health and medical needs of the high-risk youth in the juvenile justice system.

References

- 1. Broschmann, R. et. al. (2020). The Health of Adolescents in Detention: a Global Scoping Review. The Lancet Public Health, 5(2), 114-126.
- 2. Lantos, H., Wilkinson, A., Winslow, H. et al. (2019). Describing associations between child maltreatment frequency and the frequency and timing of subsequent delinquent or criminal behaviors across development: variation by sex, sexual orientation, and race. BMC Public Health 19, 1306.
- 3. Ryan, J. P., Williams, A. B., & Courtney, M. E. (2013). Adolescent neglect, juvenile delinquency and the risk of recidivism. Journal of Youth and Adolescence, 42(3), 454-465.
- 4. Yun, I., Ball, J.D., & Lim, H. (2011). Disentangling the relationship between child maltreatment and violent delinquency: Using a nationally representative sample. Journal of Interpersonal Violence, 26(1), 88–110.
- 5. Asscher, J., Van der Put, C., & Stams, G. (2015). Gender differences in the impact of abuse and neglect victimization on adolescent offending behavior. Journal of Family Violence, 30(2), 215-225.
- 6. Braga, T. et al. (2017). Unraveling the link between maltreatment and juvenile antisocial behavior: A meta-analysis of prospective longitudinal studies. Aggression and Violent Behavior (33), 37-50.
- 7. Sickmund, M., & Puzzanchera, C. (2014). Juvenile Offenders and Victims: 2014 National Report. Pittsburgh, PA: National Center for Juvenile Justice.
- 8. Skinner-Osci, et al. (2019). Justice-Involved Youth and Trauma-Informed Interventions. Justice Policy Journal, 16(2), 1-25. From http://www.cjcj.org/jpj
- 9. Acoca, L., Stephens, J., & Van Vleet, A. (2014). Health coverage and care for youth in the juvenile justice system: The Role of Medicaid and CHIP. Henry J. Kaiser Family Foundation, Washington, DC.
- 10. Leasy, M., O'Gurek, D., & Savoy, M. (2019). Unlocking Clues to Current Health and Past History: Childhood Trauma and Healing. American Academy of Family Physicians.
- 11. Kopak, A.M., & Proctor, S.L. (2016). Acute and Chronic Effects of Substance Use as Predictors of Criminal Offense Types Among Juvenile Offenders. Journal of Juvenile Justice, 5(1), 50.
- 12. Doran, N., Luczak, S.E., Bekman, N., Koutsenok, I., & Brown, S.A. (2012). Adolescent substance use and aggression a review. Criminal Justice and Behavior, 39(6), 748-769.

 13. Chassin, L., Piquero, A.R., Losoya, S.H., Mansion, A.D., & Schubert, C.A. (2013). Joint consideration of distal and proximal predictors of premature mortality among serious juvenile
- offenders. Journal of Adolescent Health, 52(6), 689-696.
 14. Lansford, J.E., Dodge, K.A., Fontaine, R.G., Bates, J.E., & Pettit, G.S. (2014). Peer rejection, affiliation with deviant peers, delinquency, and risky sexual behavior. Journal of Youth and
- Adolescence, 43(10), 1742-1751.
- 15. Winter, V.R., Brandon-Friedman, R.A., & Ely, G. (2016). Sexual health behaviors and outcomes among current and former foster youth: A review of the literature. Children and Youth Services Review. 64. 1-14.
- 16. Schuster, R.M., Mermelstein, R., & Wakschlag, L. (2013). Gender-specific relationships between depressive symptoms, marijuana use, parental communication and risky sexual behavior in adolescence. Journal of Youth and Adolescence, 42(8), 1194-1209.